4-(1,1,3,3-テトラメチルブチル)フェノールの p53KO マウスを用いた 強制経口投与による中期がん原性試験報告書

試験番号:0906,0907

FIGURES

FIGURES

(Test substance-treated group)

- FIGURE A 1 SURVIVAL RATE OF MALE p53 KO MICE IN THE 26-WEEK CARCINOGENICITY STUDY OF 4-(1,1,3,3-TETRAMETHYLBUTYL) PHENOL (GAVAGE STUDY)
- FIGURE A 2 SURVIVAL RATE OF FEMALE p53 KO MICE IN THE 26-WEEK CARCINOGENICITY STUDY OF 4-(1,1,3,3-TETRAMETHYLBUTYL) PHENOL (GAVAGE STUDY)
- FIGURE A 3 BODY WEIGHT CHANGES OF MALE p53 KO MICE IN THE 26-WEEK CARCINOGENICITY STUDY OF 4-(1,1,3,3-TETRAMETHYLBUTYL) PHENOL (GAVAGE STUDY)
- FIGURE A 4 BODY WEIGHT CHANGES OF FEMALE p53 KO MICE IN THE 26-WEEK CARCINOGENICITY STUDY OF 4-(1,1,3,3-TETRAMETHYLBUTYL) PHENOL (GAVAGE STUDY)
- FIGURE A 5 FOOD CONSUMPTION CHANGES OF MALE p53 KO MICE IN THE 26-WEEK CARCINOGENICITY STUDY OF 4-(1,1,3,3-TETRAMETHYLBUTYL) PHENOL (GAVAGE STUDY)
- FIGURE A 6 FOOD CONSUMPTION CHANGES OF FEMALE p53 KO MICE IN THE 26-WEEK CARCINOGENICITY STUDY OF 4-(1,1,3,3-TETRAMETHYLBUTYL) PHENOL (GAVAGE STUDY)

FIGURES (Positive control group)

- FIGURE B 1 SURVIVAL RATE OF MALE p53 KO MICE IN THE 26-WEEK CARCINOGENICITY STUDY OF 2-METHOXY-5-METHYLANILINE (GAVAGE STUDY)
- FIGURE B 2 SURVIVAL RATE OF FEMALE p53 KO MICE IN THE 26-WEEK CARCINOGENICITY STUDY OF 2-METHOXY-5-METHYLANILINE (GAVAGE STUDY)
- FIGURE B 3 BODY WEIGHT CHANGES OF MALE p53 KO MICE IN THE 26-WEEK CARCINOGENICITY STUDY OF 2-METHOXY-5-METHYLANILINE (GAVAGE STUDY)
- FIGURE B 4 BODY WEIGHT CHANGES OF FEMALE p53 KO MICE IN THE 26-WEEK CARCINOGENICITY STUDY OF 2-METHOXY-5-METHYLANILINE (GAVAGE STUDY)
- FIGURE B 5 FOOD CONSUMPTION CHANGES OF MALE p53 KO MICE IN THE 26-WEEK CARCINOGENICITY STUDY OF 2-METHOXY-5-METHYLANILINE (GAVAGE STUDY)
- FIGURE B 6 FOOD CONSUMPTION CHANGES OF FMALE p53 KO MICE IN THE 26-WEEK CARCINOGENICITY STUDY OF 2-METHOXY-5-METHYLANILINE (GAVAGE STUDY)




FIGURE A 1 SURVIVAL RATE OF MALE p53 KO MICE IN THE 26-WEEK CARCINOGENICITY STUDY OF 4-(1,1,3,3-TETRAMETHYLBUTYL) PHENOL (GAVAGE STUDY)




FIGURE A 2 SURVIVAL RATE OF FEMALE p53 KO MICE IN THE 26-WEEK CARCINOGENICITY STUDY OF 4-(1,1,3,3-TETRAMETHYLBUTYL) PHENOL (GAVAGE STUDY)




FIGURE A 3 BODY WEIGHT CHANGES OF MALE p53 KO MICE IN THE 26-WEEK CARCINOGENICITY STUDY OF 4-(1,1,3,3-TETRAMETHYLBUTYL) PHENOL (GAVAGE STUDY)




FIGURE A 4 BODY WEIGHT CHANGES OF FEMALE p53 KO MICE IN THE 26-WEEK CARCINOGENICITY STUDY OF 4-(1,1,3,3-TETRAMETHYLBUTYL) PHENOL (GAVAGE STUDY)




FIGURE A 5 FOOD CONSUMPTION CHANGES OF MALE p53 KO MICE IN THE 26-WEEK CARCINOGENICITY STUDY OF 4-(1,1,3,3-TETRAMETHYLBUTYL) PHENOL (GAVAGE STUDY)




FIGURE A 6 FOOD CONSUMPTION CHANGES OF FEMALE p53 KO MICE IN THE 26-WEEK CARCINOGENICITY STUDY OF 4-(1,1,3,3-TETRAMETHYLBUTYL) PHENOL (GAVAGE STUDY)




FIGURE B 1 SURVIVAL RATE OF MALE p53 KO MICE IN THE 26-WEEK CARCINOGENICITY STUDY OF 2-METHOXY-5-METHYLANILINE (GAVAGE STUDY)




FIGURE B 2 SURVIVAL RATE OF FEMALE p53 KO MICE IN THE 26-WEEK CARCINOGENICITY STUDY OF 2-METHOXY-5-METHYLANILINE (GAVAGE STUDY)




FIGURE B 3 BODY WEIGHT CHANGES OF MALE p53 KO MICE IN THE 26-WEEK CARCINOGENICITY STUDY OF 2-METHOXY-5-METHYLANILINE (GAVAGE STUDY)




FIGURE B 4 BODY WEIGHT CHANGES OF FEMALE p53 KO MICE IN THE 26-WEEK CARCINOGENICITY STUDY OF 2-METHOXY-5-METHYLANILINE (GAVAGE STUDY)




FIGURE B 5 FOOD CONSUMPTION CHANGES OF MALE p53 KO MICE IN THE 26-WEEK CARCINOGENICITY STUDY OF 2-METHOXY-5-METHYLANILINE (GAVAGE STUDY)




FIGURE B 6 FOOD CONSUMPTION CHANGES OF FEMALE p53 KO MICE IN THE 26-WEEK CARCINOGENICITY STUDY OF 2-METHOXY-5-METHYLANILINE (GAVAGE STUDY)