

1-ブロモ-3-クロロプロバンのマウスを用いた  
吸入による 2 週間毒性試験報告書

試験番号：0380

# TABLES

## TABLES

TABLE 1 SURVIVAL ANIMAL NUMBERS AND BODY WEIGHT CHANGES OF MALE MICE IN THE 2-WEEK INHALATION STUDY OF 1-BROMO-3-CHLOROPROPANE

TABLE 2 SURVIVAL ANIMAL NUMBERS AND BODY WEIGHT CHANGES OF FEMALE MICE IN THE 2-WEEK INHALATION STUDY OF 1-BROMO-3-CHLOROPROPANE

TABLE 3 FOOD CONSUMPTION CHANGES OF MALE MICE IN THE 2-WEEK INHALATION STUDY OF 1-BROMO-3-CHLOROPROPANE

TABLE 4 FOOD CONSUMPTION CHANGES OF FEMALE MICE IN THE 2-WEEK INHALATION STUDY OF 1-BROMO-3-CHLOROPROPANE

TABLE 1 SURVIVAL ANIMAL NUMBERS AND BODY WEIGHT CHANGES OF MALE MICE IN THE 2-WEEK INHALATION STUDY OF 1-BROMO-3-CHLOROPROPANE

| Week-Day on Study | 0ppm      |                    | 50ppm     |                 | 100ppm        |           | 200ppm          |               | 400ppm    |                 | 800ppm        |           |                 |               |           |     |      |
|-------------------|-----------|--------------------|-----------|-----------------|---------------|-----------|-----------------|---------------|-----------|-----------------|---------------|-----------|-----------------|---------------|-----------|-----|------|
|                   | Av.Wt.    | No.of Surviv. < 5> | Av.Wt.    | % of cont. < 5> | No.of Surviv. | Av.Wt.    | % of cont. < 5> | No.of Surviv. | Av.Wt.    | % of cont. < 5> | No.of Surviv. | Av.Wt.    | % of cont. < 5> | No.of Surviv. |           |     |      |
| 0-0               | 22.2 ( 5) | 5/ 5               | 22.1 ( 5) | 100             | 5/ 5          | 22.2 ( 5) | 100             | 5/ 5          | 22.2 ( 5) | 100             | 5/ 5          | 22.1 ( 5) | 100             | 5/ 5          | 22.1 ( 5) | 100 | 5/ 5 |
| 1-2               | 22.7 ( 5) | 5/ 5               | 22.5 ( 5) | 99              | 5/ 5          | 22.2 ( 5) | 98              | 5/ 5          | 22.2 ( 5) | 98              | 5/ 5          | 22.3 ( 5) | 98              | 5/ 5          | 21.5 ( 1) | 95  | 1/ 5 |
| 1-4               | 23.2 ( 5) | 5/ 5               | 22.7 ( 5) | 98              | 5/ 5          | 22.9 ( 5) | 99              | 5/ 5          | 22.6 ( 5) | 97              | 5/ 5          | 23.2 ( 5) | 100             | 5/ 5          | 20.7 ( 1) | 89  | 1/ 5 |
| 1-7               | 23.9 ( 5) | 5/ 5               | 23.0 ( 5) | 96              | 5/ 5          | 23.0 ( 5) | 96              | 5/ 5          | 22.7 ( 5) | 95              | 5/ 5          | 22.5 ( 5) | 94              | 5/ 5          | 21.3 ( 1) | 89  | 1/ 5 |
| 2-3               | 24.1 ( 5) | 5/ 5               | 23.5 ( 5) | 98              | 5/ 5          | 23.7 ( 5) | 98              | 5/ 5          | 23.4 ( 5) | 97              | 5/ 5          | 22.2 ( 5) | 92              | 5/ 5          | 21.4 ( 1) | 89  | 1/ 5 |
| 2-7               | 24.8 ( 5) | 5/ 5               | 23.7 ( 5) | 96              | 5/ 5          | 24.3 ( 5) | 98              | 5/ 5          | 23.8 ( 5) | 96              | 5/ 5          | 22.7 ( 5) | 92              | 5/ 5          | 21.6 ( 1) | 87  | 1/ 5 |

< > : No.of effective animals, ( ) : No.of measured animals Av.Wt. : g

TABLE 2 SURVIVAL ANIMAL NUMBERS AND BODY WEIGHT CHANGES OF FEMALE MICE IN THE 2-WEEK INHALATION STUDY OF 1-BROMO-3-CHLOROPROPANE

| Week-Day on Study | 0ppm      |                    | 50ppm     |                 | 100ppm        |           | 200ppm          |               | 400ppm    |                 | 800ppm        |           |                 |               |           |     |      |
|-------------------|-----------|--------------------|-----------|-----------------|---------------|-----------|-----------------|---------------|-----------|-----------------|---------------|-----------|-----------------|---------------|-----------|-----|------|
|                   | Av.Wt.    | No.of Surviv. < 5> | Av.Wt.    | % of cont. < 5> | No.of Surviv. | Av.Wt.    | % of cont. < 5> | No.of Surviv. | Av.Wt.    | % of cont. < 5> | No.of Surviv. | Av.Wt.    | % of cont. < 5> | No.of Surviv. |           |     |      |
| 0-0               | 18.8 ( 5) | 5/ 5               | 18.9 ( 5) | 101             | 5/ 5          | 18.9 ( 5) | 101             | 5/ 5          | 18.9 ( 5) | 101             | 5/ 5          | 18.8 ( 5) | 100             | 5/ 5          | 18.9 ( 5) | 101 | 5/ 5 |
| 1-2               | 18.8 ( 5) | 5/ 5               | 18.6 ( 5) | 99              | 5/ 5          | 19.1 ( 5) | 102             | 5/ 5          | 19.0 ( 5) | 101             | 5/ 5          | 18.7 ( 5) | 99              | 5/ 5          | 18.2 ( 2) | 97  | 2/ 5 |
| 1-4               | 19.3 ( 5) | 5/ 5               | 19.5 ( 5) | 101             | 5/ 5          | 19.6 ( 5) | 102             | 5/ 5          | 20.0 ( 5) | 104             | 5/ 5          | 19.8 ( 5) | 103             | 5/ 5          | 16.4 ( 1) | 85  | 1/ 5 |
| 1-7               | 20.1 ( 5) | 5/ 5               | 19.3 ( 5) | 96              | 5/ 5          | 19.8 ( 5) | 99              | 5/ 5          | 19.7 ( 5) | 98              | 5/ 5          | 19.3 ( 5) | 96              | 5/ 5          | 13.0 ( 1) | 65  | 1/ 5 |
| 2-3               | 20.2 ( 5) | 5/ 5               | 20.5 ( 5) | 101             | 5/ 5          | 20.9 ( 5) | 103             | 5/ 5          | 20.8 ( 5) | 103             | 5/ 5          | 19.6 ( 5) | 97              | 5/ 5          | - (-)     | -   | 0/ 5 |
| 2-7               | 20.6 ( 5) | 5/ 5               | 20.8 ( 5) | 101             | 5/ 5          | 20.6 ( 5) | 100             | 5/ 5          | 21.1 ( 5) | 102             | 5/ 5          | 19.9 ( 5) | 97              | 5/ 5          | - (-)     | -   | 0/ 5 |

< > : No.of effective animals, ( ) : No.of measured animals Av.Wt. : g

TABLE 3 FOOD CONSUMPTION CHANGES OF MALE MICE IN THE 2-WEEK INHALATION STUDY OF 1-BROMO-3-CHLOROPROPANE

| Week-Day<br>on Study | 0ppm     |                          | 50ppm    |                       | 100ppm           |          | 200ppm                |                  | 400ppm   |                       | 800ppm           |          |                       |                  |          |    |      |
|----------------------|----------|--------------------------|----------|-----------------------|------------------|----------|-----------------------|------------------|----------|-----------------------|------------------|----------|-----------------------|------------------|----------|----|------|
|                      | Av.FC.   | No.of<br>Surviv.<br>< 5> | Av.FC.   | % of<br>cont.<br>< 5> | No.of<br>Surviv. | Av.FC.   | % of<br>cont.<br>< 5> | No.of<br>Surviv. | Av.FC.   | % of<br>cont.<br>< 5> | No.of<br>Surviv. | Av.FC.   | % of<br>cont.<br>< 5> | No.of<br>Surviv. |          |    |      |
| 1-7                  | 4.6 ( 5) | 5/ 5                     | 4.5 ( 5) | 98                    | 5/ 5             | 4.3 ( 5) | 93                    | 5/ 5             | 4.2 ( 5) | 91                    | 5/ 5             | 4.1 ( 5) | 89                    | 5/ 5             | 3.0 ( 1) | 65 | 1/ 5 |
| 2-7                  | 4.3 ( 5) | 5/ 5                     | 4.3 ( 5) | 100                   | 5/ 5             | 4.4 ( 5) | 102                   | 5/ 5             | 4.4 ( 5) | 102                   | 5/ 5             | 3.9 ( 5) | 91                    | 5/ 5             | 3.6 ( 1) | 84 | 1/ 5 |

< > : No.of effective animals, ( ) : No.of measured animals      Av.FC. : g

TABLE 4 FOOD CONSUMPTION CHANGES OF FEMALE MICE IN THE 2-WEEK INHALATION STUDY OF 1-BROMO-3-CHLOROPROPANE

| Week-Day<br>on Study | 0ppm     |                          | 50ppm    |                       | 100ppm           |          | 200ppm                |                  | 400ppm   |                       | 800ppm           |          |                       |                  |          |   |      |
|----------------------|----------|--------------------------|----------|-----------------------|------------------|----------|-----------------------|------------------|----------|-----------------------|------------------|----------|-----------------------|------------------|----------|---|------|
|                      | Av.FC.   | No.of<br>Surviv.<br>< 5> | Av.FC.   | % of<br>cont.<br>< 5> | No.of<br>Surviv. | Av.FC.   | % of<br>cont.<br>< 5> | No.of<br>Surviv. | Av.FC.   | % of<br>cont.<br>< 5> | No.of<br>Surviv. | Av.FC.   | % of<br>cont.<br>< 5> | No.of<br>Surviv. |          |   |      |
| 1-7                  | 3.9 ( 5) | 5/ 5                     | 3.9 ( 5) | 100                   | 5/ 5             | 3.5 ( 5) | 90                    | 5/ 5             | 3.8 ( 5) | 97                    | 5/ 5             | 3.7 ( 5) | 95                    | 5/ 5             | 0.2 ( 1) | 5 | 1/ 5 |
| 2-7                  | 3.7 ( 5) | 5/ 5                     | 4.1 ( 5) | 111                   | 5/ 5             | 3.8 ( 5) | 103                   | 5/ 5             | 4.0 ( 5) | 108                   | 5/ 5             | 3.7 ( 5) | 100                   | 5/ 5             | - ( -)   | - | 0/ 5 |

< > : No.of effective animals, ( ) : No.of measured animals      Av.FC. : g